Ivan Friedrich, M.D. Kenneth P. Rubin, M.D. Vincent S. Panella, M.D. Mark S. Sapienza, M.D. Irina Kaplounov, M.D. Sandarsh R. Kancherla, M.D.

Eugene S. Han, M.D. Diplomates, American Board of Gastroenterology and Internal Medicine

> 420 Grand Avenue Englewood, N.J. 07631 (201)569-7044

## SuTab Split Colonoscopy Prep

	NAME:	PHYSICIAN:	
	DAY AND DATE:	TIME:	ARRIVE AT:
	PLACE:	REPORT TO:	
Ple	Please note: There is a 24 hour cancellation policy. For procedures cancelled with less than 24 hours notice, a fee of		
	25 dollars will be assessed.	•	

## 1 week Before your test

## **DAY Before your** test

## **DAY** of your test

- Check with our office for specific instructions if you take ANY blood thinning medications (Coumadin, Pradaxa, Effient, Aspirin (Ecotrin), Eliquis, Xarelto, Plavix, etc.)
- Check with your physician for specific instructions if you take ANY diabetes medications (Insulin, Metformin, Janumet, Glipizide, etc.)
- Pick up SuTab from your pharmacy.
- STOP oral iron 5 days before test (Vitamins with iron are OK)
- Do not take oral medications within 2 hours of taking SuTab doses \*
- START a CLEAR LIQUID DIET when you wake up
- NO RED LIQUIDS (No milk, No orange juice)
- Gatorade, juice, water, ice pops, tea, coffee are OK
- At **6PM** TAKE 12 tablets of SuTab over 15-20 minutes with at least 16oz of water.
- THEN at **7PM** DRINK another 16 oz of water over 30 minutes.
- At **8PM** DRINK another 16 oz of water over 30 minutes.
- Continue CLEAR LIQUID diet
- Get some sleep

• 6 hours BEFORE your test TAKE 12 tablets of SuTab over 15-20

\*

- minutes with at least 16oz of water.
- 5 hours BEFORE your test DRINK another 16 oz of water over 30
- 3 hours BEFORE your test DRINK another 16 oz of water over 30 minutes.
  - TAKE all your usual medications when you wake up with a sip of water
  - Continue CLEAR LIQUID diet
  - 2 Hours before your test: STOP ALL oral intake-NOTHING!!!

NO gum or candy

NO cologne, perfume, or lotions please!

- You must have a ride home-NO Driving until the following day!
- For all women of child-bearing age, a urine pregnancy test must be done on arrival at the center.